

Vaccinations

Babies



Birth

- Hepatitis B



2 months

- Diphtheria, tetanus, pertussis (whooping cough), hepatitis B, polio, Haemophilus influenzae type b (Hib)
- Pneumococcal
- Rotavirus



4 months

- Diphtheria, tetanus, pertussis (whooping cough), hepatitis B, polio, Haemophilus influenzae type b (Hib)
- Pneumococcal
- Rotavirus



6 months

- Diphtheria, tetanus, pertussis (whooping cough), hepatitis B, polio, Haemophilus influenzae type b (Hib)
- *Pneumococcal* *^



12 months

- Meningococcal ACWY
- Measles, mumps, rubella
- Pneumococcal
- *Hepatitis A* *



18 months

- Haemophilus influenzae type b (Hib)
- Measles, mumps, rubella, varicella (chickenpox)
- Diphtheria, tetanus, pertussis (whooping cough)
- *Hepatitis A* *



4 years

- Diphtheria, tetanus, pertussis (whooping cough), polio
- *Pneumococcal* ^

* Additional vaccines for Aboriginal and Torres Strait Islander children (QLD, NT, WA and SA). Check with your GP.

^ Additional vaccines for medically at-risk children. Check with your GP.

Pregnant women



- Pertussis (whooping cough)
- Influenza

Influenza can be given at any stage during pregnancy, but pertussis is recommended at 20-32 weeks. Check with your GP whether you need any other vaccinations.

