

TRUSAMIDE

Dorzolamide hydrochloride

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about TRUSAMIDE. It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using TRUSAMIDE against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What TRUSAMIDE is used for

TRUSAMIDE is used to lower raised pressure in the eye and to treat glaucoma. Glaucoma is a condition in which the pressure of fluid in the eye may be high. However, some people with glaucoma may have normal eye pressure. Also, some people with raised eye pressure may not have glaucoma.

Glaucoma is caused by a build-up of the fluid which flows through the eye. This build-up occurs because the fluid drains out of your eye more slowly than it is being pumped in. Since new fluid continues to enter the eye, joining the fluid already there, the pressure continues to rise. This raised pressure may damage the back of the eye resulting in gradual loss of sight. Damage can progress so slowly

that the person is not aware of this gradual loss of sight.

Sometimes even normal eye pressure is associated with damage to the back of the eye.

There are usually no symptoms of glaucoma. The only way of knowing that you have glaucoma is to have your eye pressure, optic nerve and visual field checked by an eye specialist or optometrist. If glaucoma is not treated it can lead to serious problems. You may have no symptoms but eventually glaucoma can lead to total blindness. In fact, untreated glaucoma is one of the most common causes of blindness.

Although TRUSAMIDE helps control your glaucoma it does not cure it.

For more information about glaucoma, contact Glaucoma Australia Inc., PO Box 420, Crows Nest 1585, telephone 1800 500 880.

TRUSAMIDE is used, either alone or in combination with other eye drops or medicines, to lower raised pressure within your eye(s).

TRUSAMIDE lowers pressure in the eye by reducing the production of fluid.

TRUSAMIDE belongs to a family of medicines called carbonic anhydrase inhibitors.

TRUSAMIDE is not addictive.

Before you use TRUSAMIDE

When you must not take it

Do not use TRUSAMIDE if:

- **you have an allergy to TRUSAMIDE or any of the ingredients listed at the end of this leaflet**
- **you are breast-feeding or intend to breast-feed**

It is not known whether TRUSAMIDE passes into breast milk.

Do not put the eye drops into your eye(s) while you are wearing soft contact lenses.

The preservative in TRUSAMIDE (benzalkonium chloride) may be deposited in soft contact lenses. You can put your soft contact lenses back into your eyes at least 15 minutes after you have used TRUSAMIDE.

Do not use TRUSAMIDE if:

- **the seal around the cap is broken**
- **the bottle shows signs of tampering**
- **the expiry date on the pack has passed.**

If you use this medicine after the expiry date has passed, it may not work.

If you are not sure whether you should start using TRUSAMIDE, talk to your doctor.

Do not give TRUSAMIDE to a child.

The safety and effectiveness of TRUSAMIDE in children have not been established.

Before you start to use it

Tell your doctor if:

1. **you are pregnant or intend to become pregnant**

Your doctor will discuss the risks and benefits of using TRUSAMIDE during pregnancy and a decision can be made if you should or should not use it.

2. you have now or have had in the past any medical conditions, especially the following:

- kidney disease
- liver disease

3. you have an allergy to sulfonamide medicines

The active ingredient of TRUSAMIDE, dorzolamide hydrochloride, is a sulfonamide-related compound. Therefore, if you are allergic to sulfonamide medicines you may be allergic to TRUSAMIDE. Check with your doctor or pharmacist if you are not sure whether you are allergic to sulfonamides.

4. you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes.

If you have not told your doctor about any of the above, tell them before you use TRUSAMIDE.

Taking other medicines

Tell your doctor if you are using any other medicines or eye drops, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and TRUSAMIDE may interfere with each other. These include:

- tablets used to treat glaucoma
- large amounts of aspirin or salicylates

These medicines may be affected by TRUSAMIDE, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while using TRUSAMIDE.

How to use TRUSAMIDE

How much to use

Your doctor will tell you how many drops you need to use each day.

Use TRUSAMIDE only when prescribed by your doctor.

When TRUSAMIDE is used alone, the usual dose for adults is one drop three times a day, in either one or both eyes.

If your doctor has recommended that you use TRUSAMIDE with a beta-blocker eye drop, then the usual dose for adults is one drop of TRUSAMIDE twice a day, in either one or both eyes.

Follow all directions given to you by your doctor carefully.

They may differ from the information contained in this leaflet.

After using TRUSAMIDE, wait at least 10 minutes before putting any other eye drops in your eye(s).

When to use it

If you are using TRUSAMIDE three times a day, use the drops first thing in the morning, in the early afternoon and at bedtime (i.e. approximately 8 hours apart). If you are using TRUSAMIDE twice a day, use the drops in the morning and in the evening (i.e. approximately 12 hours apart).

Use TRUSAMIDE every day, at about the same time each day, unless your doctor tells you otherwise.

Using your eye drops at the same time each day will have the best effect on your eye pressure. It will also help you remember when to use the eye drops.

How to use it

You may find it easier to put drops in your eye while you are sitting or lying down.

Before opening the bottle for the first time, make sure the two safety

seals joining the cap to the bottle are not broken. If they are, do not use the bottle and return it to your pharmacist.

You will notice a small space between the cap and the bottle - this is normal.

If you are wearing soft contact lenses, remove them before putting the drops in your eye.

1. Wash your hands well with soap and water.

2. To open the bottle for the first time, hold the bottle upright and turn the cap in the direction of the arrows until you can lift it off. This will break the two safety seals.

3. Place the cap upside down (arrows face down) on a flat surface. Do not touch the inside of the cap. This will help keep the inside of the cap clean and keep germs out of the eye drops.

4. Hold the bottle upside down in one hand, with your thumb or index finger over the "finger push" area.

5. Using your other hand, gently pull down your lower eyelid to form a pouch.

6. Tilt your head back and look up.

7. Put the tip of the bottle close to your lower eyelid. Do not let it touch your eye.

8. Release one drop into the pouch formed between your eye and eyelid by gently squeezing the bottle.

9. Close your eye and keep it closed. Do not blink or rub your eye.

10. While your eye is still closed, place your index finger against the inside corner of your eye and press against your nose for about two minutes.

This will help to stop the medicine from draining through the tear duct to the nose and throat, from where it can be absorbed into other parts of your body. Ask your doctor for more specific instructions on this technique.

11. Replace the cap, sealing it tightly. Do not overtighten the cap.

12. Wash your hands again with soap and water to remove any residue.

Wait at least 15 minutes before replacing your contact lenses.

Be careful not to touch the dropper tip against your eye, eyelid or anything else to avoid contaminating the eye drops.

Contaminated eye drops may give you an eye infection.

You may feel a slight burning sensation in the eye shortly after using the eye drops.

If this persists, or is very uncomfortable, contact your doctor or pharmacist.

How long to use it

TRUSAMIDE helps control your condition but does not cure it. Therefore, TRUSAMIDE must be used every day. Continue using TRUSAMIDE for as long as your doctor prescribes,

If you forget to use it

If it is almost time for your next dose, skip the dose you missed and use your next dose when you are meant to. Otherwise, use the drops as soon as you remember, and then go back to using them as you would normally.

If you are not sure whether to skip the dose, talk to your doctor or pharmacist.

Do not use double the amount to make up for the dose that you missed.

If you have trouble remembering to use your eye drops, ask your pharmacist for some hints.

If you use too much (overdose)

If you think that you or anyone else may have swallowed any or all of the contents of a bottle of TRUSAMIDE, or used too many drops, immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) for advice, or go to accident and

emergency at your nearest hospital. Do this even if there are no signs of discomfort or poisoning.

While you are using TRUSAMIDE

Things you must do

Have your eye pressure checked when your eye specialist says, to make sure TRUSAMIDE is working.

If you develop an eye infection, receive an eye injury, or have eye surgery tell your doctor.

Your doctor may tell you to use a new container of TRUSAMIDE because of possible contamination of the old one, or may advise you to stop your treatment with TRUSAMIDE.

If you become pregnant while using TRUSAMIDE tell your doctor.

If you are about to be started on any new medicine tell your doctor and pharmacist that you are using TRUSAMIDE.

Things you must not do

Do not give TRUSAMIDE to anyone else, even if they have the same condition as you.

Do not stop using TRUSAMIDE without first talking to your doctor.

If you stop using your eye drops, your eye pressures may rise again and damage to your eye may occur.

Things to be careful of

Be careful driving or operating machinery until you know how TRUSAMIDE affects you.

TRUSAMIDE generally does not cause any problems with your ability to drive a car or operate machinery. However, it may cause certain side effects in some people, including blurred vision and dizziness. Make

sure you know how you react to TRUSAMIDE before you drive a car or operate machinery.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using TRUSAMIDE.

TRUSAMIDE helps most people with high eye pressure and glaucoma, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- eye problems such as burning, stinging, itching, conjunctivitis, watering of the eye(s), redness of the eye(s), swelling or crusting of the eyelids(s), eye pain, blurred vision
- feeling sick
- bitter taste, dry mouth
- nose bleeds
- throat irritation
- headache, dizziness
- tiredness, weakness
- kidney stones
- numbness or tingling in fingers or toes

These are usually mild side effects of TRUSAMIDE.

If any of the following happen, stop using TRUSAMIDE and tell your doctor immediately or go to accident and emergency at your nearest hospital:

- wheezing, difficulty in breathing, shortness of breath
- swelling of the face, lips, mouth, tongue or throat which may cause

difficulty in breathing or swallowing

- severe and sudden onset of pinkish, itchy swellings on the skin, also called hives or nettlerash
- skin rash, itchiness

These may be serious side effects. You may have an allergic reaction to TRUSAMIDE. You may need urgent medical attention. Serious side effects are rare.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice any other effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

After using TRUSAMIDE

Storage

Keep your eye drops in a cool dry place where the temperature stays below 30°C.

Do not store it or any other medicine in the bathroom or near a sink.

Do not leave it in the car or on window sills.

Do not carry the eye drops in pockets of your clothes.

Heat and dampness can destroy some medicines.

Keep the eye drops away from light.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Do not leave the cap off the bottle for any length of time to avoid contaminating the eye drops.

Disposal

Write the date on the bottle when you open the eye drops and throw out any remaining solution after four weeks.

TRUSAMIDE contains a preservative which helps prevent germs growing in the solution for the first four weeks after opening the bottle. After this time there is a greater risk that the drops may become contaminated and cause an eye infection. A new bottle should be opened.

If your doctor tells you to stop using the eye drops or they have passed their expiry date, ask your pharmacist what to do with any remaining solution.

Product description

What it looks like

TRUSAMIDE comes as eye drops in a 5 mL bottle.

Ingredients

Active ingredient:

- dorzolamide hydrochloride, equivalent to dorzolamide 2%

Inactive ingredients:

- hydroxyethylcellulose
- mannitol
- sodium citrate
- sodium hydroxide
- benzalkonium chloride as preservative

Supplier

TRUSAMIDE is supplied in Australia by:

Alphapharm Pty Ltd

Level 1, 30 The Bond
30-34 Hickson Road
Millers Point NSW 2000
www.mylan.com.au

This leaflet was prepared in July 2020.

AUST R 217251

Trusamide_cmi\Jul20/00